arXiv:1609.08649v1 [math.DG] 23 Sep 2016

Projective Curvature Tensors of Second Type Almost Geodesic Mappings

Nenad O. Vesić

Abstract

We consider equitorsion second type almost geodesic mappings of a non-symmetric affine connection space in this article. Using different computational methods, we obtained some invariants of these mappings. Last generalized Thomas projective parameter and Weyl projective tensor as invariants of a second type almost geodesic mapping of a non-symmetric affine connection space are further generalized here.

1 Introduction

A lot of research papers and monographs are dedicated to developments of the theory of differential geometry [1,2,6,8–21] and its applications [3–5,7]. Einstein (see [3–5]) concluded the symmetric affine connection theory covers researches about a gravitation. The theory electromagnetism is covered by anti-symmetric parts of affine connections. The research about non-symmetric affine connected spaces is started by L. P. Eisenhart [6].

An N-dimensional manifold $\mathcal{M}_N$ endowed with a non-symmetric affine connection ∇ (affine connection coefficients L^i_{jk} and L^i_{kj} are different) is said to be *the non-symmetric affine connection space* $\mathbb{GA}_N$. Because of the previous mentioned non-symmetry of affine connection coefficients it exists symmetric and anti-symmetric part of these coefficients respectively defined as:

$$\widetilde{S}^i_{jk} = \frac{1}{2}(L^i_{jk} + L^i_{kj}) \quad \text{and} \quad \widetilde{T}^i_{jk} = \frac{1}{2}(L^i_{jk} - L^i_{kj}). \tag{1.1}$$

2010 *Math. Subj. Classification*: Primary: 53A55; Secondary: 53B05, 53C15, 53C22

Key words: almost geodesic mapping, curvature tensor, generalization, Weyl projective tensor, Thomas projective parameter

This research is supported by Ministry of Education, Science and Technological Development, Republic of Serbia, Grant No. 174012

A symmetrization and an anti-symmetrization without division by indices i and j will be denoted as $(i \dots j)$ and $[i \dots j]$ respectively.

The magnitude $\widetilde{T}^i_{jk}$ is a torsion tensor of the space $\mathbb{GA}_N$. An affine connection space $\mathbb{A}_N$ endowed with an affine connection S which coefficients coincide with the symmetric part $\widetilde{S}^i_{jk}$ of the affine connection coefficients L^i_{jk} of the space $\mathbb{GA}_N$ is said to be *the associated space* of the space $\mathbb{GA}_N$.

There are a lot of researchers interested for a development of the non-symmetric affine connection space theory. Some significant results in this subject are obtained into the papers [10, 11, 14–19, 21].

Four kinds of covariant differentiation (see [10]) with regard to an affine connection of a non-symmetric affine connection space $\mathbb{GA}_N$ are defined as:

$$a^i_{j\underset{1}{|}k} = a^i_{j,k} + L^i_{\alpha k}a^\alpha_j - L^\alpha_{jk}a^i_\alpha, \qquad a^i_{j\underset{2}{|}k} = a^i_{j,k} + L^i_{k\alpha}a^\alpha_j - L^\alpha_{kj}a^i_\alpha, \quad (1.2)$$

$$a^i_{j\underset{3}{|}k} = a^i_{j,k} + L^i_{\alpha k}a^\alpha_j - L^\alpha_{kj}a^h_\alpha, \qquad a^i_{j\underset{4}{|}k} = a^i_{j,k} + L^i_{k\alpha}a^\alpha_j - L^\alpha_{jk}a^i_\alpha, \quad (1.3)$$

for a partial derivative denoted by comma and an indexed magnitude a^i_j.

All of these covariant derivatives become restricted to a covariant derivative

$$a^i_{j;k} = a^i_{j,k} + \widetilde{S}^i_{\alpha k}a^\alpha_j - \widetilde{S}^\alpha_{jk}a^i_\alpha, \quad (1.4)$$

of the magnitude a^i_j with regard to an affine connection of the associated space $\mathbb{A}_N$ of the space $\mathbb{GA}_N$.

For this reason, it exists only one curvature tensor

$$R^i_{jmn} = \widetilde{S}^i_{jm;n} - \widetilde{S}^i_{jn;m}, \quad (1.5)$$

of the associated space $\mathbb{A}_N$.

1.1 Almost geodesic mappings of a space $\mathbb{GA}_N$

In an attempt to generalize the term of geodesics N. S. Sinyukov (see [12]) defined an almost geodesic line of a symmetric affine connection space $\mathbb{A}_N$. Consequently, he defined a term of an almost geodesic mapping f between symmetric affine connection spaces $\mathbb{A}_N$ and $\overline{\mathbb{A}}_N$. Sinyukov noticed three types π_1, π_2, π_3 of almost geodesic mappings between symmetric affine connection spaces. His research has been directly developed by many authors in a lot of papers [1, 2, 8, 9, 13, 20].

The Sinyukov's generalization of geodesics is primary developed for the case of a generalized affine connection space $\mathbb{GA}_N$ in [14–16]. In this space it exists four kinds of covariant differentiation but these covariant derivatives are reduced onto first two ones (1.2) for the case of any contra-variant tensor.

For this reason, there are two kinds of almost geodesic lines of the space $\mathbb{GA}_N$ [14–19, 21] defined as a curve $\ell = \ell(t)$ which tangential vector $\lambda^i = d\ell^i/dt \neq 0$ satisfies the following equations:

$$\underset{\theta}{\overline{\lambda}}{}^i_{(2)} = \underset{\theta}{\overline{a}}(t)\lambda^i + \underset{\theta}{\overline{b}}(t)\underset{\theta}{\overline{\lambda}}{}^i_{(1)}, \quad \underset{\theta}{\overline{\lambda}}{}^i_{(1)} = \lambda^i_{\underset{\theta}{\|}\alpha}\lambda^\alpha, \quad \underset{\theta}{\overline{\lambda}}{}^i_{(2)} = \underset{\theta}{\overline{\lambda}}{}^i_{(1)\underset{\theta}{\|}\alpha}\lambda^\alpha, \tag{1.6}$$

$\theta = 1, 2$, for covariant differentiation of the θ-th kind with regard to affine connection of the space $\mathbb{G}\overline{\mathbb{A}}_N$ denoted by $\underset{\theta}{\|}$.

Because of two kinds of almost geodesic lines of this space a mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ is **the almost geodesic mapping of a θ-th kind**, $\theta = 1, 2$, if any geodesic line of the space $\mathbb{GA}_N$ it turns into an almost geodesic line of the θ-th kind of the space $\mathbb{G}\overline{\mathbb{A}}_N$. For this reason, there are three types of almost geodesic mappings of the space $\mathbb{GA}_N$ and any of these three types have two kinds. A class of almost geodesic mappings of a τ-th type, $\tau = 1, 2, 3$, and of a θ-th kind $\theta = 1, 2$ of the space $\mathbb{GA}_N$ is denoted as $\underset{\theta}{\pi}_\tau$.

Basic equations of a second type almost geodesic mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ of a θ-th kind, $\theta = 1, 2$, are [15]:

$$\overline{L}^i_{jk} = L^i_{jk} + \psi_j\delta^i_k + \psi_k\delta^i_j + \sigma_j F^i_k + \sigma_k F^i_j + \xi^i_{jk}, \tag{1.7}$$

$$\begin{aligned} &F^i_{j\underset{\theta}{|}k} + F^i_{k\underset{\theta}{|}j} + F^i_\alpha F^\alpha_j\sigma_k + F^i_\alpha F^\alpha_k\sigma_j + (-1)^\theta\big(\xi^i_{j\alpha}F^\alpha_k + \xi^i_{k\alpha}F^\alpha_j\big) \\ &\quad = \mu_j F^i_k + \mu_k F^i_j + \nu_j\delta^i_k + \nu_k\delta^i_j, \end{aligned} \tag{1.8}$$

for covariant vectors μ_j, ν_j, an affinor F^i_j and an anti-symmetric tensor ξ^i_{jk}.

A second type almost geodesic mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ of a θ-th kind, $\theta = 1, 2$, satisfies the property of reciprocity $\big($it is an element of the class $\underset{\theta}{\pi}_2(e)\big)$ if it saves the affinor F^i_j and its inverse mapping is a second type almost geodesic mapping of the θ-th kind. An almost geodesic mapping f of the space $\mathbb{GA}_N$ satisfies the property of reciprocity (see [15]) if and only if the affinor F^i_j satisfies a relation

$$F^i_\alpha F^\alpha_j = e\delta^i_j, \qquad e = 0, \pm 1. \tag{1.9}$$

2 Invariants of second type almost geodesic mappings

The aim of this paper is to find some new invariants of almost geodesic mappings of a second type which satisfy the property of reciprocity. The results in this subject obtained until now are about the theories of special subclasses of the classes $\underset{\theta}{\pi}_2, \theta = 1, 2$.

Motivated by Sinyukov's results, it is obtained (see [15]) magnitudes

$$\underset{1}{T}{}^i_{jk} = \widetilde{S}^i_{jk} - \frac{1}{e - F^2}\Big(\big(F\widetilde{S}^\alpha_{k\alpha} - F^\alpha_k\widetilde{S}^\beta_{\alpha\beta}\big)F^i_j + \big(F\widetilde{S}^\alpha_{j\alpha} - F^\alpha_j\widetilde{S}^\beta_{\alpha\beta}\big)F^i_k\Big), \quad (2.10)$$

$$\begin{aligned}\underset{2}{\hat{T}}{}^i_{jk} &= T^i_{jk} + eF^i_\alpha\big(\underset{1}{F}{}^\alpha_{(j|k)} - \widetilde{T}^\alpha_{\beta(k}F^\beta_{j)}\big)\\ &\quad - \frac{e}{1+N}F^\beta_\alpha\Big(\big(\underset{1}{F}{}^\alpha_{\beta|j} - \widetilde{T}^\alpha_{\gamma(\beta}F^\gamma_{j)}\big)\delta^i_k + \big(\underset{1}{F}{}^\alpha_{\beta|k} - \widetilde{T}^\alpha_{\gamma(\beta}F^\gamma_{k)}\big)\delta^i_j\Big),\end{aligned} \quad (2.11)$$

for $F = F^\alpha_\alpha$, $e - F^2 \neq 0$ and Thomas projective parameter T^i_{jk} of the associated space $\mathbb{A}_N$ in the expression of the invariant $\underset{1}{T}{}^i_{jk}$, are invariants of a canonical second type almost geodesic mapping of the first kind of the space $\mathbb{GA}_N$.

Moreover, Weyl projective tensor of the space $\mathbb{G}\hat{\mathbb{A}}_N$ which affine connection coefficients are $\hat{L}^i_{jk} = L^i_{jk} + eF^i_\alpha\underset{1}{F}{}^\alpha_{(j|k)} - eT^\alpha_{\beta(j}F^\beta_{k)}F^i_\alpha$ is an invariant of the canonical second type almost geodesic mapping f of the first kind. The aim of our following research is to find some other more general invariants of special second type almost geodesic mappings of the space $\mathbb{GA}_N$.

Let a mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion second type almost geodesic mapping of a θ-th kind, $\theta = 1, 2$, which satisfies the property of reciprocity. The composition (1.9) involved into the basic equation (1.8) together with using of the fact the mapping f is an equitorsion one ($\xi^i_{jk} = 0$) involved into the both of basic equations (1.7, 1.8) proves it is satisfied relations

$$\overline{L}^i_{jk} = L^i_{jk} + \psi_j\delta^i_k + \psi_k\delta^i_j + \sigma_jF^i_k + \sigma_kF^i_j, \quad (2.12)$$

$$\underset{\theta}{F}{}^i_{j|k} + \underset{\theta}{F}{}^i_{k|j} = \mu_jF^i_k + \mu_kF^i_j + \big(\nu_j - e\sigma_j\big)\delta^i_k + \big(\nu_k - e\sigma_k\big)\delta^i_j. \quad (2.13)$$

It is proved a following proposition is satisfied in this way.

Proposition 2.1 *Let $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion second type almost geodesic mapping of a θ-th kind, $\theta = 1, 2$, which satisfies the property of reciprocity. The equations* (2.12, 2.13, 1.9) *are basic equations of this mapping.* □

Based on the fact the second type almost geodesic mapping f satisfies the property of reciprocity the corresponding magnitudes $\overline{\psi}_i, \overline{\sigma}_i, \overline{F}^i_j$ which determine an inverse mapping f^{-1} of the mapping f are [15]

$$\overline{\psi}_i = -\psi_i, \quad \overline{\sigma}_i = -\sigma_i, \quad \overline{F}^i_j = F^i_j.$$

After contracting the basic equation (2.12) by indices i and k and using the fact it is satisfied a relation $\sigma_j = \frac{1}{2}(\sigma_j - \overline{\sigma}_j)$ we obtain it is satisfied an equation

$$\psi_j = \frac{1}{N+1}\big(\overline{L}^\alpha_{j\alpha} - L^\alpha_{j\alpha}\big) + \frac{1}{2(N+1)}\Big[\big(\overline{\sigma}_j\overline{F} + \overline{\sigma}_\alpha\overline{F}^\alpha_j\big) - \big(\sigma_j F + \sigma_\alpha F^\alpha_j\big)\Big], \quad (2.14)$$

for $F = F^\alpha_\alpha$ as above.

Using the previous expression of the magnitude ψ_j we conclude the basic equation (2.12) has a form

$$\overline{L}^i_{jk} = L^i_{jk} + \overline{\omega}^i_{jk} - \omega^i_{jk}, \quad (2.15)$$

for

$$\begin{aligned}\omega^i_{jk} = &-\frac{1}{2}(\sigma_j F^i_k + \sigma_k F^i_j) + \frac{1}{N+1}\big(L^\alpha_{j\alpha}\delta^i_k + L^\alpha_{k\alpha}\delta^i_j\big)\\ &+ \frac{1}{2(N+1)}\Big(\big(\sigma_j F + \sigma_\alpha F^\alpha_j\big)\delta^i_k + \big(\sigma_k F + \sigma_\alpha F^\alpha_k\big)\delta^i_j\Big),\end{aligned} \quad (2.16)$$

and the magnitude $\overline{\psi}_j$ defined in the same manner as a function of the corresponding elements of the space $\mathbb{G}\overline{\mathbb{A}}_N$.

The equation (2.15) proves it is satisfied an equality

$$\overline{\underset{2}{\mathcal{T}}}{}^i_{jk} = \underset{2}{\mathcal{T}}{}^i_{jk},$$

for

$$\underset{2}{\mathcal{T}}{}^i_{jk} = L^i_{jk} - \omega^i_{jk} \quad \text{and} \quad \overline{\underset{2}{\mathcal{T}}}{}^i_{jk} = \overline{L}^i_{jk} - \overline{\omega}^i_{jk}. \quad (2.17)$$

It is proved a following lemma in this way.

Lemma 2.1 *Let $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion almost geodesic mapping of a second type which satisfies the property of reciprocity. A magnitude $\underset{2}{\mathcal{T}}{}^i_{jk}$ defined in the first of the expressions* (2.17) *is an invariant of the mapping f.* □

The invariant $\underset{2}{\mathcal{T}}{}^i_{jk}$ of a second type almost geodesic mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ which satisfies the property of reciprocity is said to be **the π_2-generalized Thomas projective parameter**.

Let us generalize Weyl projective tensor of an equitorsion almost geodesic mapping $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ of the first type which satisfies the property of reciprocity. This generalization will be realized just for first type almost geodesic mappings. A result for the case of second type almost geodesic mappings may be obtained in the same manner.

First of all, we can observe the magnitude ω^i_{jk} defined into the equation (2.16) is symmetric by indices j and k. For this reason, symmetric parts $\widetilde{S}^i_{jk}$ and $\overline{\widetilde{S}}{}^i_{jk}$ of affine connection coefficients L^i_{jk} and $\overline{L}{}^i_{jk}$ of the spaces $\mathbb{GA}_N$ and $\mathbb{G}\overline{\mathbb{A}}_N$ satisfy a relation

$$\overline{\widetilde{S}}{}^i_{jk} = \widetilde{S}^i_{jk} + \overline{\omega}^i_{jk} - \omega^i_{jk}, \tag{2.18}$$

for the above defined magnitudes ω^i_{jk} and $\overline{\omega}^i_{jk}$.

Using the covariant derivative of the first kind (1.2) we obtain a curvature tensor R^i_{jmn} of the associated space $\mathbb{A}_N$ has a form:

$$\begin{aligned} R^i_{jmn} = {} & \widetilde{S}^i_{jm\underset{1}{|}n} - \widetilde{S}^i_{jn\underset{1}{|}m} - \widetilde{T}^i_{\alpha n}\widetilde{S}^\alpha_{jm} - \widetilde{T}^\alpha_{jm}\widetilde{S}^i_{\alpha n} \\ & + \widetilde{T}^\alpha_{jn}\widetilde{S}^i_{\alpha m} + \widetilde{T}^i_{\alpha m}\widetilde{S}^\alpha_{jn} + 2\widetilde{T}^\alpha_{mn}\widetilde{S}^i_{j\alpha}. \end{aligned} \tag{2.19}$$

Motivated by this result we are going to find a rule of change of the curvature tensor R^i_{jmn} bellow. Let us involve following substitutions:

$$\underset{2}{\mathcal{U}}{}^i_{jk} = \frac{1}{2}\big(\underset{2}{\mathcal{T}}{}^i_{jk} + \underset{2}{\mathcal{T}}{}^i_{kj}\big) \quad \text{and} \quad \underset{2}{\overline{\mathcal{U}}}{}^i_{jk} = \frac{1}{2}\big(\underset{2}{\overline{\mathcal{T}}}{}^i_{jk} + \underset{2}{\overline{\mathcal{T}}}{}^i_{kj}\big), \tag{2.20}$$

for the above obtained invariant $\underset{2}{\mathcal{T}}{}^i_{jk} = \underset{2}{\overline{\mathcal{T}}}{}^i_{jk}$ of the mapping f.

From the equations (2.17) and (2.20) together with the above mentioned symmetry of the magnitude ω^i_{jk} from the equation (2.16) by indices j and k we conclude it is satisfied equalities

$$\underset{2}{\mathcal{U}}{}^i_{jk} = \widetilde{S}^i_{jk} - \omega^i_{jk} \quad \text{and} \quad \underset{2}{\overline{\mathcal{U}}}{}^i_{jk} = \overline{\widetilde{S}}{}^i_{jk} - \overline{\omega}^i_{jk}. \tag{2.21}$$

It is easy to be obtained covariant derivatives

$$\underset{2}{\mathcal{U}}{}^i_{jm|n} = \frac{1}{2}\big(\underset{2}{\mathcal{T}}{}^i_{jm|n} + \underset{2}{\mathcal{T}}{}^i_{mj|n}\big) \quad \text{and} \quad \underset{2}{\overline{\mathcal{U}}}{}^i_{jm\|n} = \frac{1}{2}\big(\underset{2}{\overline{\mathcal{T}}}{}^i_{jm\|n} + \underset{2}{\overline{\mathcal{T}}}{}^i_{mj\|n}\big), \tag{2.22}$$

of the previous defined magnitudes $\underset{2}{\mathcal{U}}{}^i_{jm}$ and $\underset{2}{\overline{\mathcal{U}}}{}^i_{jm}$ satisfy a relation

$$\begin{aligned}\underset{2}{\overline{\mathcal{U}}}{}^i_{jm\|n} =& \underset{2}{\mathcal{U}}{}^i_{jm|n} + \overline{\widetilde{S}}{}^i_{\alpha n}\underset{2}{\overline{\mathcal{U}}}{}^\alpha_{jm} - \overline{\widetilde{S}}{}^\alpha_{jn}\underset{2}{\overline{\mathcal{U}}}{}^i_{\alpha m} - \overline{\widetilde{S}}{}^\alpha_{mn}\underset{2}{\overline{\mathcal{U}}}{}^i_{j\alpha} \\ & - \widetilde{S}^i_{\alpha n}\underset{2}{\mathcal{U}}{}^\alpha_{jm} + \widetilde{S}^\alpha_{jn}\underset{2}{\mathcal{U}}{}^i_{\alpha m} + \widetilde{S}^\alpha_{mn}\underset{2}{\mathcal{U}}{}^i_{j\alpha}.\end{aligned} \tag{2.23}$$

The equations (2.21, 2.23) prove it is satisfied a following proposition.

Proposition 2.2 *Let $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion second type almost geodesic mapping of the first kind between non-symmetric affine connection spaces $\mathbb{GA}_N$ and $\mathbb{G}\overline{\mathbb{A}}_N$. Covariant derivatives $\widetilde{S}^i_{jm|n}$ and $\overline{\widetilde{S}}{}^i_{jm\|n}$ of symmetric parts $\widetilde{S}^i_{jm}$ and $\overline{\widetilde{S}}{}^i_{jm}$ of the corresponding affine connection coefficients L^i_{jm} and $\overline{L}^i_{jm}$ satisfy a relation*

$$\begin{aligned}\overline{\widetilde{S}}{}^i_{jm\|n} =& \widetilde{S}^i_{jm|n} + \overline{\omega}^i_{jm\|n} - \omega^i_{jm|n} + \overline{\widetilde{S}}{}^i_{\alpha n}\underset{2}{\overline{\mathcal{U}}}{}^\alpha_{jm} - \overline{\widetilde{S}}{}^\alpha_{jn}\underset{2}{\overline{\mathcal{U}}}{}^i_{\alpha m} - \overline{\widetilde{S}}{}^\alpha_{mn}\underset{2}{\overline{\mathcal{U}}}{}^i_{j\alpha} \\ & - \widetilde{S}^i_{\alpha n}\underset{2}{\mathcal{U}}{}^\alpha_{jm} + \widetilde{S}^\alpha_{jn}\underset{2}{\mathcal{U}}{}^i_{\alpha m} + \widetilde{S}^\alpha_{mn}\underset{2}{\mathcal{U}}{}^i_{j\alpha},\end{aligned} \tag{2.24}$$

for the magnitudes $\underset{2}{\mathcal{U}}{}^i_{jk}$ and $\underset{2}{\overline{\mathcal{U}}}{}^i_{jk}$ defined above. □

Using the invariance $\overline{\widetilde{T}}{}^i_{jk} = \widetilde{T}^i_{jk}$ and the consequent invariances $\underset{2}{\overline{\mathcal{T}}}{}^\alpha_{jm}\overline{\widetilde{T}}{}^i_{\alpha n} = \underset{2}{\mathcal{T}}{}^\alpha_{jm}\widetilde{T}^i_{\alpha n}$ such as $\underset{2}{\overline{\mathcal{T}}}{}^i_{j\alpha}\overline{\widetilde{T}}{}^\alpha_{mn} = \underset{2}{\mathcal{T}}{}^i_{j\alpha}\widetilde{T}^\alpha_{mn}$ we obtain it is satisfied a following proposition.

Proposition 2.3 *Let $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion second type almost geodesic mapping of the first kind between non-symmetric affine connection spaces $\mathbb{GA}_N$ and $\mathbb{G}\overline{\mathbb{A}}_N$. Magnitudes $\widetilde{S}^\alpha_{jm}\widetilde{T}^i_{\alpha n}, \widetilde{S}^i_{\alpha j}\widetilde{T}^\alpha_{mn}$ and its deformations $\overline{\widetilde{S}}{}^\alpha_{jm}\overline{\widetilde{T}}{}^i_{\alpha n}, \overline{\widetilde{S}}{}^i_{\alpha j}\overline{\widetilde{T}}{}^\alpha_{mn}$ under the mapping f satisfy equations*

$$\widetilde{\overline{S}}^{\alpha}_{jm}\widetilde{\overline{T}}^{i}_{\alpha n} = \widetilde{S}^{\alpha}_{jm}\widetilde{T}^{i}_{\alpha n} + \overline{\omega}^{\alpha}_{jm}\widetilde{\overline{T}}^{i}_{\alpha n} - \omega^{\alpha}_{jm}\widetilde{T}^{i}_{\alpha n}, \tag{2.25}$$

$$\widetilde{\overline{S}}^{i}_{j\alpha}\widetilde{\overline{T}}^{\alpha}_{mn} = \widetilde{S}^{i}_{j\alpha}\widetilde{T}^{\alpha}_{mn} + \overline{\omega}^{i}_{j\alpha}\widetilde{\overline{T}}^{\alpha}_{mn} - \omega^{i}_{j\alpha}\widetilde{T}^{\alpha}_{mn}, \tag{2.26}$$

for the magnitude ω^{i}_{jk} *defined into the equation* (2.16) *and the corresponding one* $\overline{\omega}^{i}_{jk}$. □

If we contract the basic equation (2.13) by the indices i and k we conclude it is satisfied a relation

$$F_{\underset{\theta}{|j}} = \mu_j F + \mu_\alpha F^{\alpha}_{j} + (N+1)(\nu_j - e\sigma_j) - F^{\alpha}_{j\underset{\theta}{|\alpha}}, \tag{2.27}$$

$\theta = 1, 2$. This equation proves it is satisfied a following proposition.

Proposition 2.4 *Let* $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ *be an equitorsion second type almost geodesic mapping of the first kind between non-symmetric affine connection spaces* $\mathbb{GA}_N$ *and* $\mathbb{G}\overline{\mathbb{A}}_N$ *which satisfies the property of reciprocity. A covariant derivative* $\omega^{i}_{jm\underset{1}{|}n}$ *of the magnitude* ω^{i}_{jk} *defined into the equation* (2.16) *satisfies an equality*

$$\begin{aligned}
\omega^{i}_{jm\underset{1}{|}n} &= \frac{1}{N+1}\big(L^{\alpha}_{j\alpha\underset{1}{|}n}\delta^{i}_{m} + L^{\alpha}_{m\alpha\underset{1}{|}n}\delta^{i}_{j}\big) + \frac{1}{2}(\nu_n - e\sigma_n)(\sigma_j\delta^{i}_{m} + \sigma_m\delta^{i}_{j}) \\
&- \frac{1}{2}\big(\sigma_{j\underset{1}{|}n}F^{i}_{m} + \sigma_{m\underset{1}{|}n}F^{i}_{j} + \sigma_j F^{i}_{m\underset{1}{|}n} + \sigma_m F^{i}_{j\underset{1}{|}n}\big) \\
&+ \frac{1}{2(N+1)}\big(\sigma_{j\underset{1}{|}n}F + \sigma_{\alpha\underset{1}{|}n}F^{\alpha}_{j} + \sigma_\alpha F^{\alpha}_{j\underset{1}{|}n}\big)\delta^{i}_{m} \\
&+ \frac{1}{2(N+1)}\big(\sigma_{m\underset{1}{|}n}F + \sigma_{\alpha\underset{1}{|}n}F^{\alpha}_{m} + \sigma_\alpha F^{\alpha}_{m\underset{1}{|}n}\big)\delta^{i}_{j} \\
&+ \frac{1}{2(N+1)}\big(\mu_n F + \mu_\alpha F^{\alpha}_{n} - F^{\alpha}_{n\underset{1}{|}\alpha}\big)(\sigma_j\delta^{i}_{m} + \sigma_m\delta^{i}_{j}),
\end{aligned} \tag{2.28}$$

for magnitudes μ_i, ν_i *used into the basic equation* (2.13). □

A difference $\widehat{\Delta}^{i}_{jmn} = \overline{\omega}^{i}_{jm\underset{1}{\|}n} - \omega^{i}_{jm\underset{1}{|}n}$, of the magnitudes $\overline{\omega}^{i}_{jm\underset{1}{\|}n}$ and $\omega^{i}_{jm\underset{1}{|}n}$ satisfies a relation

$$\begin{aligned}
\widehat{\Delta}^{i}_{jmn} &= \frac{1}{N+1}\Big(\big(\widetilde{\overline{S}}^{\alpha}_{j\alpha\underset{1}{\|}n} - \widetilde{S}^{\alpha}_{j\alpha\underset{1}{|}n}\big)\delta^{i}_{m} + \big(\widetilde{\overline{S}}^{\alpha}_{m\alpha\underset{1}{\|}n} - \widetilde{S}^{\alpha}_{m\alpha\underset{1}{|}n}\big)\delta^{i}_{j}\Big) \\
&+ \hat{\overline{\rho}}^{i}_{jmn} - \hat{\rho}^{i}_{jmn},
\end{aligned} \tag{2.29}$$

for

$$
\begin{aligned}
2\hat{\rho}^i_{jmn} &= -2L^\beta_{jn}\widetilde{T}^\alpha_{\beta\alpha}\delta^i_m - 2L^\beta_{mn}\widetilde{T}^\alpha_{\beta\alpha}\delta^i_j + (\nu_n - e\sigma_n)(\sigma_j\delta^i_m + \sigma_m\delta^i_j)\\
&- (\sigma_{j\underset{1}{|}n}F^i_m + \sigma_{m\underset{1}{|}n}F^i_j + \sigma_j F^i_{m\underset{1}{|}n} + \sigma_m F^i_{j\underset{1}{|}n})\\
&+ \frac{1}{N+1}\big(\sigma_{j\underset{1}{|}n}F + \sigma_{\alpha\underset{1}{|}n}F^\alpha_j + \sigma_\alpha F^\alpha_{j\underset{1}{|}n}\big)\delta^i_m\\
&+ \frac{1}{N+1}\big(\sigma_{m\underset{1}{|}n}F + \sigma_{\alpha\underset{1}{|}n}F^\alpha_m + \sigma_\alpha F^\alpha_{m\underset{1}{|}n}\big)\delta^i_j\\
&+ \frac{1}{N+1}\big(\mu_n F + \mu_\alpha F^\alpha_n - F^\alpha_{n\underset{1}{|}\alpha}\big)(\sigma_j\delta^i_m + \sigma_m\delta^i_j),
\end{aligned}
\tag{2.30}
$$

a magnitude $\hat{\overline{\rho}}^i_{jmn}$ from the space $\mathbb{G}\overline{\mathbb{A}}_N$ analogue to the magnitude $\hat{\rho}^i_{jmn}$, the magnitudes μ_i, ν_i from the equation (2.13) and the corresponding ones $\overline{\mu}_i, \overline{\nu}_i$.

From the equations (2.29, 2.30) we conclude it exists a magnitude

$$
\begin{aligned}
\hat{\upsilon}_{\underset{1}{ij}} &= -\frac{1}{N+1}\widetilde{\overline{S}}^\alpha_{i\alpha\underset{1}{\|}j} + \overline{L}^\beta_{ij}\widetilde{\overline{T}}^\alpha_{\beta\alpha} - (\overline{\nu}_j - e\overline{\sigma}_j)\sigma_i\\
&+ \frac{1}{2(N+1)}\big(\overline{\sigma}_{i\underset{1}{\|}j}\overline{F} + \overline{\sigma}_{\alpha\underset{1}{\|}j}\overline{F}^\alpha_i + \overline{\sigma}_\alpha\overline{F}^\alpha_{i\underset{1}{\|}j}\big)\\
&- \frac{1}{2(N+1)}\big(\overline{\mu}_j\overline{F} + \overline{\mu}_\alpha\overline{F}^\alpha_j - \overline{F}^\alpha_{j\underset{1}{\|}\alpha}\big)\overline{\sigma}_i\\
&+ \frac{1}{N+1}\widetilde{S}^\alpha_{i\alpha\underset{1}{|}j} - L^\beta_{ij}\widetilde{T}^\alpha_{\beta\alpha} + (\nu_j - e\sigma_j)\sigma_i\\
&- \frac{1}{2(N+1)}\big(\sigma_{i\underset{1}{|}j}F + \sigma_{\alpha\underset{1}{|}j}F^\alpha_i + \sigma_\alpha F^\alpha_{i\underset{1}{|}j}\big)\\
&+ \frac{1}{2(N+1)}\big(\mu_j F + \mu_\alpha F^\alpha_j - F^\alpha_{j\underset{1}{|}\alpha}\big)\sigma_i,
\end{aligned}
\tag{2.31}
$$

such that the equation (2.24) has a form

$$
\begin{aligned}
\widetilde{\overline{S}}^i_{jm\underset{1}{\|}n} &= \widetilde{S}^i_{jm\underset{1}{|}n} - \delta^i_m\hat{\upsilon}_{\underset{1}{jn}} - \delta^i_j\hat{\upsilon}_{\underset{1}{mn}}\\
&- \frac{1}{2}\big(\overline{\sigma}_{j\underset{1}{\|}n}\overline{F}^i_m + \overline{\sigma}_{m\underset{1}{\|}n}\overline{F}^i_j + \overline{\sigma}_j\overline{F}^i_{m\underset{1}{\|}n} + \overline{\sigma}_m\overline{F}^i_{j\underset{1}{\|}n}\big)\\
&+ \frac{1}{2}\big(\sigma_{j\underset{1}{|}n}F^i_m + \sigma_{m\underset{1}{|}n}F^i_j + \sigma_j F^i_{m\underset{1}{|}n} + \sigma_m F^i_{j\underset{1}{|}n}\big)\\
&+ \widetilde{\overline{S}}^i_{\alpha n}\overline{\mathcal{U}}^\alpha_{\underset{2}{jm}} - \widetilde{\overline{S}}^\alpha_{jn}\overline{\mathcal{U}}^i_{\underset{2}{\alpha m}} - \widetilde{\overline{S}}^\alpha_{mn}\overline{\mathcal{U}}^i_{\underset{2}{j\alpha}} - \widetilde{S}^i_{\alpha n}\mathcal{U}^\alpha_{\underset{2}{jm}} + \widetilde{S}^\alpha_{jn}\mathcal{U}^i_{\underset{2}{\alpha m}} + \widetilde{S}^\alpha_{mn}\mathcal{U}^i_{\underset{2}{j\alpha}}.
\end{aligned}
\tag{2.32}
$$

Using the equations (2.20, 2.25, 2.26, 2.32) and the expression (2.19) of the curvature tensors R^i_{jmn} and $\overline{R}^i_{jmn}$ of the associated spaces $\mathbb{A}_N$ and $\overline{\mathbb{A}}_N$ we obtain it is satisfied the following equation

$$\begin{aligned}
\overline{R}^i_{jmn} &= R^i_{jmn} - \delta^i_m \underset{1}{\hat{v}}_{jn} + \delta^i_n \underset{1}{\hat{v}}_{jm} - \delta^i_j \underset{1}{\hat{v}}_{[mn]} \\
&- \frac{1}{2}\big(\overline{\sigma}_{j\underset{1}{\|}n}\overline{F}^i_m + \overline{\sigma}_{m\underset{1}{\|}n}\overline{F}^i_j + \overline{\sigma}_j\overline{F}^i_{m\underset{1}{\|}n} + \overline{\sigma}_m\overline{F}^i_{j\underset{1}{\|}n}\big) \\
&+ \frac{1}{2}\big(\sigma_{j\underset{1}{|}n}F^i_m + \sigma_{m\underset{1}{|}n}F^i_j + \sigma_j F^i_{m\underset{1}{|}n} + \sigma_m F^i_{j\underset{1}{|}n}\big) \\
&+ \frac{1}{2}\big(\overline{\sigma}_{j\underset{1}{\|}m}\overline{F}^i_n + \overline{\sigma}_{n\underset{1}{\|}m}\overline{F}^i_j + \overline{\sigma}_j\overline{F}^i_{n\underset{1}{\|}m} + \overline{\sigma}_n\overline{F}^i_{j\underset{1}{\|}m}\big) \\
&- \frac{1}{2}\big(\sigma_{j\underset{1}{|}m}F^i_n + \sigma_{n\underset{1}{|}m}F^i_j + \sigma_j F^i_{n\underset{1}{|}m} + \sigma_n F^i_{j\underset{1}{|}m}\big) \\
&+ 2\widetilde{\overline{S}}^i_{\alpha n}\widetilde{\overline{S}}^\alpha_{jm} - 2\widetilde{\overline{S}}^\alpha_{jn}\widetilde{\overline{S}}^i_{\alpha m} - 2\widetilde{S}^i_{\alpha n}\widetilde{S}^\alpha_{jm} + 2\widetilde{S}^\alpha_{jn}\widetilde{S}^i_{\alpha m} \\
&- \overline{\omega}^\alpha_{jm}\overline{L}^i_{\alpha n} + \omega^\alpha_{jm}L^i_{\alpha n} - \overline{\omega}^i_{\alpha n}\overline{L}^\alpha_{jm} + \omega^i_{\alpha n}L^\alpha_{jm} \\
&+ \overline{\omega}^i_{\alpha m}\overline{L}^\alpha_{jn} - \omega^i_{\alpha m}L^\alpha_{jn} + \overline{\omega}^\alpha_{jn}\overline{L}^i_{\alpha m} - \omega^\alpha_{jn}L^i_{\alpha m} + 2\overline{\omega}^i_{j\alpha}\widetilde{\overline{T}}^\alpha_{mn} - 2\omega^i_{j\alpha}\widetilde{T}^\alpha_{mn} \\
&= R^i_{jmn} - \delta^i_m \underset{1}{\hat{v}}_{jn} + \delta^i_n \underset{1}{\hat{v}}_{jm} - \delta^i_j \underset{1}{\hat{v}}_{[mn]} + \underset{1}{\mathcal{F}}{}^i_{jmn} - \underset{1}{\overline{\mathcal{F}}}{}^i_{jmn},
\end{aligned} \tag{2.33}$$

for

$$\begin{aligned}
\underset{1}{\mathcal{F}}{}^i_{jmn} &= \frac{1}{2}\big(\sigma_{j\underset{1}{|}n}F^i_m + \sigma_{m\underset{1}{|}n}F^i_j + \sigma_j F^i_{m\underset{1}{|}n} + \sigma_m F^i_{j\underset{1}{|}n}\big) \\
&- \frac{1}{2}\big(\sigma_{j\underset{1}{|}m}F^i_n + \sigma_{n\underset{1}{|}m}F^i_j + \sigma_j F^i_{n\underset{1}{|}m} + \sigma_n F^i_{j\underset{1}{|}m}\big) \\
&- 2\widetilde{S}^i_{\alpha n}\widetilde{S}^\alpha_{jm} + 2\widetilde{S}^\alpha_{jn}\widetilde{S}^i_{\alpha m} \\
&+ \omega^\alpha_{jm}L^i_{\alpha n} + \omega^i_{\alpha n}L^\alpha_{jm} - \omega^i_{\alpha m}L^\alpha_{jn} - \omega^\alpha_{jn}L^i_{\alpha m} - 2\omega^i_{j\alpha}\widetilde{T}^\alpha_{mn},
\end{aligned} \tag{2.34}$$

the magnitude ω^i_{jk} defined into the equation (2.16) and the corresponding magnitude $\underset{1}{\overline{\mathcal{F}}}{}^i_{jmn}$.

After contracting the equation (2.33) by indices i and n we conclude Ricci tensors R_{jm} and $\overline{R}_{jm}$ of the associated spaces $\mathbb{A}_N$ and $\overline{\mathbb{A}}_N$ satisfy a relation

$$\overline{R}_{jm} = R_{jm} + (N-1)\underset{1}{\hat{v}}_{jm} + \underset{1}{\hat{v}}_{[jm]} + \underset{1}{\mathcal{F}}_{jm} - \underset{1}{\overline{\mathcal{F}}}_{jm}, \tag{2.35}$$

for $\underset{1}{\mathcal{F}}_{jm} = \underset{1}{\mathcal{F}}{}^\alpha_{jm\alpha}$ and $\underset{1}{\overline{\mathcal{F}}}_{jm} = \underset{1}{\overline{\mathcal{F}}}{}^\alpha_{jm\alpha}$.

After alternating the equation (2.35) by indices j and m we conclude it is satisfied a relation

$$(N+1)\underset{1}{\hat{v}}_{[jm]} = \overline{R}_{[jm]} - R_{[jm]} - \underset{1}{\mathcal{F}}_{[jm]} + \underset{1}{\overline{\mathcal{F}}}_{[jm]}. \tag{2.36}$$

From the equations (2.35) and (2.36) we conclude it is satisfied an expression

$$\begin{aligned}(N^2-1)\underset{1}{\hat{v}}_{jm} = (N\overline{R}_{jm} + \overline{R}_{mj}) - (NR_{jm} + R_{mj})\\ + (N\underset{1}{\overline{\mathcal{F}}}_{jm} + \underset{1}{\overline{\mathcal{F}}}_{mj}) - (N\underset{1}{\mathcal{F}}_{jm} + \underset{1}{\mathcal{F}}_{mj}).\end{aligned} \tag{2.37}$$

After involving the results (2.36, 2.37) in the equation (2.33) we obtain it is satisfied an equality

$$\underset{2}{\overline{\mathcal{W}}}^{i}_{jmn} = \underset{2}{\mathcal{W}}^{i}_{jmn},$$

where we denoted by

$$\begin{aligned}\underset{2}{\mathcal{W}}^{i}_{jmn} = R^{i}_{jmn} + \frac{1}{N+1}\delta^{i}_{j}R_{[mn]} + \frac{N}{N^2-1}\delta^{i}_{[m}R_{jn]} + \frac{1}{N^2-1}\delta^{i}_{[m}R_{n]j}\\ + \underset{1}{\mathcal{F}}^{i}_{jmn} + \frac{1}{N+1}\delta^{i}_{j}\underset{1}{\mathcal{F}}_{[mn]} + \frac{N}{N^2-1}\delta^{i}_{[m}\underset{1}{\mathcal{F}}_{jn]} + \frac{1}{N^2-1}\delta^{i}_{[m}\underset{1}{\mathcal{F}}_{n]j},\end{aligned} \tag{2.38}$$

a geometric object of the space $\mathbb{GA}_N$, where the magnitude $\underset{1}{\mathcal{F}}^{i}_{jmn}$ is defined into the equation (2.34) and for the corresponding magnitude $\underset{1}{\mathcal{F}}_{ij} = \underset{1}{\mathcal{F}}^{\alpha}_{ij\alpha}$. The corresponding magnitude $\underset{2}{\overline{\mathcal{W}}}^{i}_{jmn}$ of the space $\mathbb{G}\overline{\mathbb{A}}_N$ is defined in the same manner.

It is proved a following theorem is satisfied in this way.

Theorem 2.1 *Let $f : \mathbb{GA}_N \to \mathbb{G}\overline{\mathbb{A}}_N$ be an equitorsion second type almost geodesic mapping of the first kind which satisfies the property of reciprocity. The magnitude $\underset{2}{\mathcal{W}}^{i}_{jmn}$ defined into the equation* (2.38) *is an invariant of this mapping.* □

3 Acknowledgements

This paper is financially supported by the Serbian Ministry of Education, Science and Technological Developments, Grant. No. 174012.

References

[1] V. Berezovski, J. Mikeš, *On the classification of almost geodesic mappings of affine-connected spaces*, Differential geometry and its applications (Dubrovnik, 1988), 41-48, Univ. Novi Sad, Novi Sad, 1989, MR 1040054.

[2] V. E. Berezovski, J. Mikeš, A. Vanžurová, *Fundamental PDE's of the Canonical Almost Geodesic Mappings of Type* $\widetilde{\pi}_1$, Bull. Malays. Math. Sci. Soc. (2) 37, No. 3, 647-659 (2014). Zbl 1298.53018.

[3] A. Einstein, *A Generalization of the Relativistic Theory of Gravitation*, Annals of Mathematics, Princeton, 46, (1945), 576-584.

[4] A. Einstein, *Bianchi Identities in the Generalized Theory of Gravitation*, Canadian Journal of Mathematics, 2, (1950), 120-128.

[5] A. Einstein, *Relativistic Theory of the Non-symmetric field*, Appendix II in the Book: The Meaning of Relativity 5^{th} edit., Princeton, 49, 1955.

[6] L. P. Eisenhart, *Non-Riemannian Geometry*, American Mathematical Society, Colloquium Publications, vol. VIII, New York, 1927.

[7] S. Ivanov, M. Zlatanović, *Connections on Non-symmetric (Generalized) Riemannian Manifold and Gravity*, Classical and Quantum Gravity, Volume 33, Number 7, 075016, (2016).

[8] J. Mikeš, H. Chudá, I. Hinterleitner, *Conformal holomorphically projective mappings of almost Hermitian manifolds with a certain initial condition*, Int. J. Geom. Methods Mod. Phys. 11, No. 5, Article ID 1450044, 8 p. (2014), Zbl 1304.53012.

[9] J. Mikeš, at all, *Differential geometry of special mappings*, Olomouc: Palacký University, Faculty of Science (ISBN 978-80-244-4671-4/pbk). 566 p. (2015), Zbl 06556646.

[10] S. M. Minčić, *Ricci identities in the space of non-symmetric affine connection*, Matematički Vesinik, 10(25), Vol. 2, (1973), 161-172.

[11] S. M. Minčić, *Independent curvature tensors and pseudotensors of spaces with non-symmetric affine connexion*, Colloquia Mathematica Societatis Janos Bolayai, 31, Differential Geometry, Budapest (Hungary), (1979), 445-460.

[12] N. S. Sinyukov, *Geodesic mappings of Riemannian spaces*, "Nauka" Moskow, (1979) (in Russian).

[13] V. S. Sobchuk, J. Mikeš, O. Pokorná, *On Almost Geodesic Mappings π_2 between Semisymmetric Riemannian Spaces*, Novi Sad J. Math., Vol. 29, No. 3, 1999, 309-312.

[14] M. S. Stanković, *First type almost geodesic mappings of general affine connection spaces*, Novi Sad J. Math. 29 (3) (1999) 313-323.

[15] M. S. Stanković, *On a canonic almost geodesic mappings of the second type of affine spaces*, FILOMAT 13 (1999) 105-114.

[16] M. S. Stanković, *On a special almost geodesic mappings of the third type of affine spaces*, Novi Sad J. Math. 31 (2) (2001) 125-135.

[17] M. S. Stanković, *Special equitorsion almost geodesic mappings of the third type of non-symmetric affine connection spaces*, Applied Mathematics and Computation, 244, (2014), 695-701.

[18] M. S. Stanković, N. O. Vesić, *Some relations in non-symmetric affine connection spaces with regard to a special almost geodesic mappings of the third type*, FILOMAT 29:9 (2015), 1941-1951.

[19] M. S. Stanković, M. Lj. Zlatanović, N. O. Vesić, *Basic Equations of G-Almost Geodesic Mappings of the Second Type, which Have the Property of Reciprocity*, CZECH MATH J, 65 (140) (2015), 787-799.

[20] H. Vavříková, J. Mikeš, O. Pokorná, G. Starko, *On fundamental equations of almost geodesic mappings of type $\pi_2(e)$*, Russ. Math. vol. 51 (1) (2007) 8-12.

[21] N. O. Vesić, *Curvature Tensors and the Third Type Almost Geodesic Mappings*, FACTA UNIVERSITATIS (NIŠ), Ser. Math. Inform. Vol. 29, No 4 (2014), 445-460.

Author's addresses:

- *Nenad O. Vesić*;

Address: Faculty of Sciences and Mathematics,
Višegradska 33, 18000 Niš, Serbia;
e-mail: vesko1985@pmf.ni.ac.rs